

CONFIRMATION OF OCCURRENCE OF THE NARROW-CLAWED CRAYFISH ? *STACUS LEPTODACTYLUS* ESCHSCHOLTZ, 1823 IN THE RIVER EVROS IN GREECE

C. PERDIKARIS (1), E. KOUTRAKIS (2), V. SARAGLIDOU (3) and N. MARGARIS (4)

-
- (1) Technological Educational Institute of Epirus, Department of Aquaculture & Fisheries, Irinis & Filias 1, GR-46100 Igoumenitsa. Greece
E-Mail: kperdik@otenet.gr
- (2) Fisheries Research Institute of Kavala, National Agricultural Research Foundation, GR-64007, Nea Peramos, Kavala, Greece.
E-Mail: manosk@inale.gr
- (3) Prefecture of Rodopi-Evros, County of Northern Evros, Fisheries Department, GR-68200, Orestiada, Greece.
- (4) Department of Environmental Studies, University of the Aegean, University Hill, GR-81100, Mytilene, Greece.
E-Mail: margaris@dolnet.gr

Reçu le 8 février 2007
Accepté le 1^{er} juin 2007

Received February 8, 2007
Accepted June 1, 2007

ABSTRACT

The narrow-clawed crayfish *Astacus leptodactylus* Eschscholtz, 1823, is widely distributed in eastern European and western Asian countries. However, species presence in Greece was documented only by limited museum conserved specimens dating back to 1893. Recently (May 2005) several crayfish were captured in the Evros River (close to city of Didimotihó) and these were identified as *A. leptodactylus*. The present work confirms the presence of *A. leptodactylus* in the Evros River still hundred and twelve years later and introduces the possibility of *A. leptodactylus* being indigenous to Greece.

Key-words: narrow-clawed crayfish, *Astacus leptodactylus*, Evros River, Greece.

PRÉSENCE DE L'ÉCREVISSE À PIEDS GRÊLES ? *STACUS LEPTODACTYLUS* ESCHSCHOLTZ, 1823 EN GRÈCE DANS LE FLEUVE EVROS

RÉSUMÉ

L'écrevisse à pieds grêles *Astacus leptodactylus* Eschscholtz, 1823, est largement répandue en Europe de l'Est et dans de nombreux pays de l'Ouest de l'Asie. Cependant, la présence de cette espèce n'est documentée en Grèce que par des spécimens conservés dans des collections de musée, remontant à 1893. Récemment (mai 2005) plusieurs écrevisses ont été capturées dans le fleuve Evros (près de la ville de Didimotihó), et identifiées comme *A. leptodactylus*. Le présent travail confirme la présence de *A. leptodactylus* dans le fleuve Evros cent douze ans plus tard et présente la possibilité que *A. leptodactylus* est indigène en Grèce.

Mots-clés: écrevisse à pieds grêles, *Astacus leptodactylus*, fleuve Evros, Grèce.

INTRODUCTION

The narrow-clawed crayfish *Astacus leptodactylus* Eschscholtz, 1823 is one of the five species of the Astacidae family endemic to the European continent. Its original distribution extends to the Ponto-Caspian region (HOLDICH, 2002; MACHINO and HOLDICH, 2006; HOLDICH *et al.*, 2006). Additionally, it has been imported to several European Union countries (SKURDAL and TAUGBØL, 2002; MACHINO and HOLDICH, 2006), mainly for farming purposes or replacing *Astacus astacus* (Linnaeus 1758) stocks devastated by the crayfish plague (*Aphanomyces astaci* Schikora, 1906).

Early reports on the species presence in Turkey were published by NINNI (1923: p. 61; under names of *Astacus fluviatilis*, *Astacus pallipes* and *Astacus leptodactylus*), BOTT (1950), TORTONESE (1952: p. 84; under name of *Potamobius leptodactylus*), HOLTHUIS (1961), KARAMAN (1962, 1963), GELDIAY and KOCATAS (1970), and PRETZMANN (1973). Then ROTH and KINZELBACH (1986), KÖKSAL (1988) and HOLDICH *et al.* (2006) mapped the distribution of the species after sample collections and previous bibliographical sources. For many years, huge amounts of live *A. leptodactylus* were harvested in Turkish lakes (ROTH and KINZELBACH, 1986; KÖKSAL, 1988; HARLIOĞLU, 2004; annual harvest and export figures cited in KÖKSAL, 1988 and HOLDICH and WHISSON, 2004). The crayfish were exported live, deep frozen or cooked mainly to Swedish, French and German markets. Indigenous stocks of *A. leptodactylus* seem to be in a recovery phase after a serious decline in the populations due to the overfishing and crayfish plague. The total commercial catches of crayfish in Turkey declined from 5,000 tons in 1984 to 200 tons in 1991, while in 2000 the total annual catches increased to 1,600-1,900 tons (HOLDICH *et al.*, 2006).

Populations of *A. leptodactylus* exist also in Bulgaria (BULGURKOV, 1961; SUBCHEV and STANIMIROVA, 1998): many sites across the Danube system and other waters of the Black Sea drainage. The species distribution has been expanded, mainly due to the water quality improvement, which is probably due to the fact that many industries have suspended their activities (ZIKOV, 1999). Nevertheless there has been no report about the existence of the species in the Bulgarian part of the River Evros (Maritsa, Meriç) yet.

The fact that *A. leptodactylus* is present in both Bulgaria and Turkey raised questions about possible current occurrence in north-eastern Greece (Evros Prefecture in Region of East Macedonia and Thrace), especially in the system of the Evros River.

Bibliographical sources on *A. leptodactylus* occurrence in Greek waters are extremely scarce. KARAMAN (1963) and KÖKSAL (1988) mentioned that *A. leptodactylus* was found in the Evros River system. But historical presence of *A. leptodactylus* in Greece has only been documented by few samples collected from the Evros River dating back to 19th century and held in the Naturhistorisches Museum in Wien (NHW) (MACHINO and HOLDICH, 2006): one specimen from the Evros River close to the city of Soufli (3 October 1893, inventory No. 3090); one specimen from a tributary of the River Evros close to the Mandra village (October 1893, inventory No. 3093); and four specimens from the cities of Alexandroupolis close the delta of the River Evros and Edirne (former Adrianoupolis) on the Turkish side (inventory No. 2873) (Figure 1). KARAMAN (1963) observed these specimens and knew that *A. leptodactylus* lived in the Evros River on both sides, Greek and Turkish. He treated the species as indigenous there. But, as he did not specify the exact places on the Greek territory, later astacologists of today did not remark or even did ignore what KARAMAN (1963) meant about presence of *A. leptodactylus* on the Greek side of the Evros drainage. Thus till recent time *A. leptodactylus* had been regarded as unknown and exotic to Greece. For example, endemism of the species to Greece was denied by KALLISTORATOS (1990) and WESTMAN *et al.* (1990). Nevertheless, occasionally the species was assigned as "native to Greece" as well, e.g., by SKURDAL and TAUGBØL (2002). Unfortunately the latter authors failed to develop their argument

about *A. leptodactylus* as native to Greece. Based on the crayfish collections in the Vienna museum (NHW), MACHINO and HOLDICH (2006) suggested that species be present in Greece and probably indigenous to.

After collected materials, the present work provides for the first time after 1893 a proof of *A. leptodactylus* occurrence in the Greek part of the Evros River.




Figure 1


Map of the north-eastern part of Greece where the Evros River outflows to the Aegean Sea. Black circle indicates where the specimens of *A. leptodactylus* were caught in May 2005, while the open circles indicate where the museum specimens are coming from (based on the description reported by MACHINO and HOLDICH, 2006). The places located in the figure are not the exact locations, but the most probable, according to the explanations given in the text.

Figure 1

Carte de la région nord-est de la Grèce où le fleuve Evros se jette dans la mer Égée. Le cercle noir indique le lieu où des spécimens de *A. leptodactylus* ont été capturés en mai 2005, alors que les cercles blancs montrent les endroits d'où proviennent les spécimens de musée (basé sur la description rapportée par MACHINO et HOLDICH, 2006). Les lieux indiqués ne représentent pas les lieux exacts, mais les plus probables, selon les explications indiquées dans le texte.

METHODS AND RESULTS

In order to verify information about the occurrence of crayfish presence in the area of Didimotiho, May 5, 2006, two fyke-net traps (80 cm diameter open end and mesh size of 6 mm) were placed on two sites in the Evros River close to the city of Didimotiho (Figure 1) with the collaboration of a local fisherman. Both sites were in the main body of the river and traps were placed at a depth of 3 m without bait. The river substrate on both sites was sandy and with rich vegetation (aquatic macrophytes and tree roots). The following day, six crayfish were found trapped; one adult male specimen was preserved in 10% formaldehyde solution (Figure 2 a, b) while the five others were released unharmed. All the specimens were identified as *A. leptodactylus* according to criteria referred in KARAMAN (1963), ALBRECHT (1982), SKURDAL and TAUGBØL (2002), FÜREDER and MACHINO (2002) and HOLDICH and VIGNEUX (2006). The conserved male specimen was 10.41 cm in total length and had characteristic narrow and thin chela. Description, sex and morphological characteristics of the specimen are shown in Table I.



Figures 2a, b

Dorsal (a) and ventral (b) view of the male specimen of *A. leptodactylus* caught close to the city of Didimotiho in the River Evros.

Figure 2a, b

Vue dorsale (a) et ventrale (b) du spécimen mâle de *A. leptodactylus* capturé près de la ville de Didimotiho dans le fleuve Evros.

The Evros basin, including the Arda, Tundja and Ergene tributaries of the lower basin, is one of the major river systems located in the eastern Balkans, with a total length of 550 km (the second longest river on the Balkan Peninsula after the Danube) and a total catchment area of 39,000 km². About 66% belongs to Bulgaria, 28% to Turkey and 6% to Greece. About 218 km of the river are located in Greece, with 203 km of the river forming the borderline with Turkey. The catchment area in Greece is 3,340 km² (INWEB, 2006).

Table I

Description, sex and morphological characteristics of *A. leptodactylus* specimen caught in the River Evros in May 2005.

Tableau I

Description, sexe et caractéristiques morphologiques du spécimen de *A. leptodactylus* capturé dans le fleuve Evros en mai 2005.

Sex	Male
Cephalothorax (mm)	52.56
Rostrum (mm)	14.55
Acumen (mm)	6.97 (pointed with two spines on each lateral margin of the rostrum)
Tail without telson (mm)	36.61
Telson (mm)	14.96
Eye diameter (mm)	3.41
“Talon” (pleopod 2)	present
Chela	narrow and long
Rostrum	elongated
Rostral borders	almost parallel, denticulated
Rostral median crista	present (14.27 mm long) with spines
Post-orbital ridges	two pairs (1 st pair immediately behind the eyes). Each ridge with one sharp spine on the front
Carapace	very “rough” with four lateral spines behind of each side of the cervical groove (one pair of spines very prominent)
Distal border of maxilliped 3 merus	one prominent hard spine on the outer front part (close to carpus)
Chela surface granulation	big
Colouration	dark patches dorsally to the carapace, pereopods and claws and pale coloration ventrally to the body

The Evros River in Greece has poor water quality since high concentrations of nitrogen and phosphorus were measured (SKOULIKIDIS *et al.*, 1998) and its annual dissolved inorganic nitrogen (DIN) and dissolved inorganic phosphorus (DIP) loads were found to be the highest after comparison with other Greek rivers (SYLAIOS *et al.*, 2005) as it receives transborder pollution. The river suffers from flow disturbances (i.e. damming, dyking and fragmentation). Moreover, during the last decade severe floods have occurred along the Evros River and its tributaries and caused a lot of damage to downstream areas. For example, the alert and alarm water level for floods coming from the Evros River are set to 4.70 and 5.70 m respectively and the actual level reached 6.92 m March 14, 2006, while the normal water levels are less than 3 m (official unpublished data from the Department of Political Emergency Planning, Prefecture of Evros). Flooding is mainly observed during springtime with heavy rainfall and snowmelt. It is also reinforced by the overflowing of artificial lakes in the Bulgarian part of the Ardas River (tributary of the Evros River).

DISCUSSION

The identified *A. leptodactylus* specimens confirmed that the species lives in the Evros River and forms one or several populations. Moreover, the Evros represents one of the south-western limits of the species' natural distribution (HOLDICH *et al.*, 2006) and probably a unique habitat of the species in Greece. Old data (i.e., museum collections), coupled with the present findings, suggest that the species could be indigenous in Greece.

Among crayfish species in Europe, those of the genus *Astacus* are of the Ponto-Caspian origin, while those of the genus *Austropotamobius* are rather of western origin (the Balkans and western Europe) (HOBBS, 1988; HOLDICH *et al.*, 2006). Looking at the crayfish distribution maps of Europe (HOLDICH *et al.*, 2006), occurrence of *A. leptodactylus* in Evros River is not an isolated point within its distribution area. After the biogeographic pattern around the Ponto-Caspian basin, the distribution in the Evros clearly appears non-scattered from the main distribution. Based on biogeographical data for fish fauna (ECONOMIDIS and BANARESCU, 1991), it can be suggested that *A. leptodactylus* have reached the Evros River system along with other Ponto-Caspian taxa of brackish water or slow-running freshwater origin through the freshwater discharges of Black Sea to the Aegean Sea. The dispersal root from the north through the Stara Planina Mountain (i.e. rheophilic species of Danubian origin) should be excluded for *A. leptodactylus*, as it is not rheophilic species and is characterised by wide ecological tolerance (e.g. salinity). Recently other crayfish populations were reportedly located by local fishermen in the Pithio trench close to the city of Orestiada (Evros Prefecture, Region of East Macedonia and Thrace) and in the Ardas River (Figure 1). The vicinity of places (Didimotihó vis-à-vis the Pithio trench and the Ardas River) strongly suggests that these crayfish probably belong to the crayfish species of Didimotihó (i.e., *A. leptodactylus*). Nevertheless occurrence of another crayfish *Austropotamobius torrentium* (Schränk, 1803) in the Bulgarian parts of the Evros drainage (BULGURKOV, 1961; SUBCHEV and STANIMIROVA, 1998) and in the River Velika (tributary of the River Rezovska) in European Turkey (TRONTELJ *et al.*, 2005) suggest that there be a possibility that *A. torrentium* is present in the Greek parts of the Evros drainage. But the main body of the Evros as well as the two crayfish places in the Ardas River and Pithio trench is characterised by muddy substrate without habitats for *A. torrentium*, thus *A. torrentium* occurrence at the two place is certainly excluded.

Apart from the Evros River system, the deliberate release or accidental escape of specimens in other freshwater or estuarine ecosystems cannot be excluded. Greece imported 4.00, 3.17 and 0.54 tons of live *A. leptodactylus* during 1983, 1984, and 1985, respectively (KÖKSAL, 1988). For example, several specimens may have been released into the River Pinios drainage in central Greece during the 1980s (PALEOCOSTAS pers. comm.), although this should be verified on the field yet.

The authors are in close cooperation with the local Fisheries Department in order to help making the local fishermen sensitive for the species conservation. Crayfish are considered as by-catch by the fishermen and frequently returned back to the river. But for establishing distribution maps of *A. leptodactylus* in the Evros drainage and conserving these precious populations as well as their habitats, further research and interstate actions are urgently needed. The issuing of conservation measures for the Evros River should also be considered.

ACKNOWLEDGEMENTS

The authors would like to thank Christos Goudinadis for his valuable help in crayfish trapping and Yoichi Machino for reviewing the manuscript.

REFERENCES

- ALBRECHT H., 1982. Das System der europäischen Flußkrebse (Decapoda, Astacidae): Vorschlag und Begründung. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut*, 79, 187-210.
- BOTT R., 1950. Die Flußkrebse Europas (Decapoda: Astacidae). *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 483, 1-36 + 6 pl.
- BULGURKOV K., 1961. Sistematika, biologiya i zoogeografsko razprostranenie na sladkovodnite raci ot sem. Astacidae i sem. Potamonidae v Bylgariya. *Izvestija na Zoologiceskijata Institut s Muzej*, 10, 165-192.

- ECONOMIDIS P., BANARESCU P.M., 1991. The distribution and origins of freshwater fishes in the Balkan Peninsula, especially in Greece. *Internationale Revue der Gesamten Hydrobiologie*, 76 (2), 257-284.
- FÜREDER L., MACHINO Y., 2002. A revised key of freshwater crayfish in Europe. *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck*, 89, 169-178 + Erratum.
- GELDIAY R., KOCATAS A., 1970. The preliminary report about the taxonomy and distribution of *Astacus* (Decapoda) in Turkey. *Scientific Reports of the Faculty of Science, Ege University*, 94, 1-7 + 3 plates, Izmir.
- HARLIOĞLU M.M., 2004. The present situation of freshwater crayfish, *Astacus leptodactylus* (Eschscholtz, 1823) in Turkey. *Aquaculture*, 230, 181-187.
- HOBBS H.H.Jr., 1988. Crayfish distribution, adaptive radiation and evolution. *In: Freshwater crayfish: Biology, management and exploitation*, HOLDICH D.M., LOWERY R.S. (eds.). Croom Helm, London, pp. 52-82 + 426-479.
- HOLDICH D.M., 2002. Present distribution of crayfish in Europe and some adjoining countries. *Bulletin Français de la Pêche et de la Pisciculture*, 367, 611-650.
- HOLDICH D.M., WHISSON G., 2004. The First 30 Years. *A history of the International Association of Astacology*. International Association of Astacology, 248 p.
- HOLDICH D.M., VIGNEUX E., 2006. Key to crayfish in Europe. *In: Atlas of crayfish in Europe*, SOUTY-GROSSET C., HOLDICH D.M., NOËL P., REYNOLDS J.D., HAFFNER P. (eds.). *Collection Patrimoines Naturels*, 64, 53-55.
- HOLDICH D.M., CARRAL J., FÜREDER L., GHERARDI F., MACHINO Y., PÖCKL M., SMIETANA P., TAUGBØL T., 2006. Species description. *In: Atlas of crayfish in Europe*, SOUTY-GROSSET C., HOLDICH D.M., NOËL P., REYNOLDS J.D., HAFFNER P. (eds.). *Collection Patrimoines Naturels*, 64, 56-129 + 163-177.
- HOLTHUIS L.B., 1961. Report on a collection of Crustacea Decapoda and Stomatopoda from Turkey and the Balkans. *Zoologische Verhandelingen*, 47, 1-67, Leiden.
- INWEB, 2006. International Network of Water-Environment Centres for the Balkans (<http://www.inweb.gr/>: 12 December 2006).
- KALLISTORATOS G., 1990. Greece. *In: Status of crayfish stocks, fisheries, diseases and culture in Europe*, WESTMAN K., PURSIANEN M., WESTMAN P. (eds.). *Kalutkimuksia-Fiskundersökningar (Helsinki)*, 3, 103-107.
- KÖKSAL G., 1988. *Astacus leptodactylus* in Europe. *In: Freshwater Crayfish: Biology, Management and Exploitation*, HOLDICH D.M., LOWERY R.S. (eds.). Croom Helm, London, pp. 365-400 + 426-479.
- KARAMAN M.S., 1962. Ein Beitrag zur Systematik des Astacidae (Decapoda). *Crustaceana*, 3 (3), 173-191.
- KARAMAN M.S., 1963. Studie der Astacidae (Crustacea, Decapoda). *Hydrobiologia (The Hague)*, 22, 111-132.
- MACHINO Y., HOLDICH D.M., 2006. Distribution of crayfish in Europe and adjoining countries: updates and comments. *Freshwater Crayfish*, 15, 292-323.
- NINNI E., 1923. Primo contributo allo studio dei pescie della pesca nelle acque dell'Imperio Ottomano. *Missione Italiana per l'Esplorazione dei Mari di Levante (materiali raccolti durante la campagna talassografica. 1921-1922 a bordo della R. Nave L.F. Marsigli)*, 5, 1-187 + 16 pl. + 1 map. Venezia.
- PRETZMANN G., 1973. Ergebnisse einiger Sammelreisen nach Vorderasien. *Annalen des Naturhistorischen Museums in Wien*, 77, 331-333.

- ROTH J., KINZELBACH R., 1986. The distribution of the Pontian Crayfish, *Astacus leptodactylus*, in Turkey. *Zoology in the Middle East*, 1, 147-152.
- SKOULIKIDIS N.T., BERTAHAS I., KOUSSOURIS T., 1998. The environmental state of freshwater resources in Greece (river and lakes). *Environmental Geology*, 36 (1/2), 1-17.
- SKURDAL J., TAUGBØL T., 2002. Crayfish of commercial importance. *Astacus*. In: *Biology of Freshwater Crayfish*, HOLDICH D.M., (ed.), pp. 467-510 + pl. 8 + pl. 19 + pl. 34-41. Blackwell Science Ltd, Oxford, UK.
- SUBCHEV M., STANIMIROVA L., 1998. Razprostranenie na pravite sladkovodni raci (Crustacea: Decapoda) i tehnite epibionti ot rod *Branchiobdella* (Annelida: Branchiobdellae), *Hystricosoma chappuisi* Michaelsen, 1926 (Annelida: Oligochaeta) i *Nitocrella divaricata* (Crustacea: Copepoda) v Bylgarija. *Historia Naturalis Bulgarica*, 9, 5-18.
- SYLAIOS G., KAMIDIS N., MARKOU D., KOUTRAKIS E., KALLIANIOTIS A., 2005. Environmental monitoring in Nestos River Delta. *Proceedings of the 12th Hellenic Congress of Ichthyologists*. Drama, Greece 13-16 October 2005, p. 258-261.
- TORTONESE E., 1952. Relazione preliminare di un viaggio a scopo zoologico attraverso l'Asia Minore. *Bollettino dell'Istituto e Museo di Zoologia della Università di Torino*, 3, 81-97 + 1 pl., Torino.
- TRONTELJ P., MACHINO Y., SKET B., 2005. Phylogenetic and phylogeographic relationships in the crayfish genus *Austropotamobius* inferred from mitochondrial COI gene sequences. *Molecular Phylogenetics and Evolution*, 34, 212-226.
- WESTMAN K., PURSIAINEN M., WESTMAN P., 1990. The situation of crayfish stocks, fisheries, diseases and crayfish culture in Europe - a review. In: *Status of crayfish stocks, fisheries, diseases and culture in Europe*, WESTMAN K., PURSIAINEN M., WESTMAN P., (eds.). *Kalaturkimuksia-Fiskundersökningar (Helsinki)*, 3, 5-31.
- ZIKOV A., 1999. Crayfish culture in Bulgaria. *Crayfish News - IAA Newsletter*, April 1999, 21 (2), 25-26.