

Feeding ecology of two newt species (*Triturus cristatus* and *Lissotriton vulgaris*) during the reproduction season

I. Roșca⁽¹⁾, I. Gherghel^{(2),*}, A. Strugariu⁽³⁾, Ș.R. Zamfirescu⁽³⁾

Received November 24, 2012

Revised January 17, 2013

Accepted February 5, 2013

ABSTRACT

Key-words:
feeding relations, resource partitioning, trophic niche, interspecific competition, Salamandridae

The aim of this study was to provide an in-depth survey of feeding ecology and trophic interactions of two syntopic newt species (*Triturus cristatus* and *Lissotriton vulgaris*) inhabiting aquatic breeding habitats from the eastern Romanian Carpathian Mountains. We sampled 736 individuals from both species. The trophic spectrum was based mostly on Asselidae (>30%). Our results show that both species may be considered generalists because their niche breadth is higher than 0.5, with largely overlapping trophic niches (>70%), which may indicate food competition.

RÉSUMÉ

L'écologie alimentaire de deux espèces de tritons (*Triturus cristatus* et *Lissotriton vulgaris*) pendant la saison de reproduction

Mots-clés :
relations alimentaires, partitionnement des ressources, niche trophique, concurrence interspécifique, Salamandridae

Le but de cette étude était d'analyser en profondeur l'écologie alimentaire et les interactions trophiques de deux espèces syntopiques de tritons (*Triturus cristatus* et *Lissotriton vulgaris*) vivant dans les habitats aquatiques dans l'est des Carpates roumaines. Nous avons échantillonné 736 individus des deux espèces. Le spectre trophique a été basé principalement sur les Asselidae (>30 %). Nos résultats montrent que les deux espèces peuvent être considérées comme des généralistes, car leur largeur de niche est supérieure à 0,5, en grande partie avec chevauchement des niches trophiques (>70 %), ce qui pourrait indiquer une concurrence alimentaire.

Amphibians are representatives for the current global biodiversity decline (Semlitsch, 2003; Stuart *et al.*, 2004). Many amphibian species require both terrestrial and aquatic habitats during their life cycle, which makes them particularly vulnerable to a wide range of detrimental factors (Alford *et al.*, 2001; Semlitsch, 2003). Newts (Order Caudata) occupy a variety of aquatic and terrestrial habitat types where they act as top-predators (Schabetsberger and Jersabek, 1995). Therefore, studies focused on their feeding ecology are necessary for

(1) Centre of Advanced Research in Bionanoconjugates and Biopolymers, "Petru Poni" Institute of Macromolecular Chemistry, 41A Aleea Grigore Ghica-Voda, 700487 Iasi, Romania

(2) Department of Zoology, Oklahoma State University, 501 Life Sciences West, 74078 Stillwater, Oklahoma, USA

(3) Faculty of Biology, "Alexandru Ioan Cuza" University, 20A Carol I Blvd., 700505 Iași, Romania

* Corresponding author: iulian.gherghel@okstate.edu

a better understanding of freshwater trophic interactions as well as for future conservation measures (Hodar, 1996).

The great crested newt (*Triturus cristatus*) and the smooth newt (*Lissotriton vulgaris*) are two widely spread Eurasian species, living in various habitats (e.g. Cogălniceanu et al., 2000). The two species differ in body size: *T. cristatus* is a large newt (up to 18 cm) and *L. vulgaris* is small to medium (up to 11–12 cm; Cogălniceanu et al., 2000). Although during the past decade many studies discussed the trophic ecology of newts (e.g. Covaciu-Marcov et al. 2010, Cicort-Lucaciu et al., 2005), most of them reported only dietary descriptions, whereas trophic interactions remained unapproached.

Here we aimed to quantify trophic niche characteristics of *T. cristatus* and *L. vulgaris*, with special emphasis on the niche breadth and niche overlap. Additionally, based on new information regarding diet preferences of the studied species, we assessed the ecological relationships between *T. cristatus* and *L. vulgaris* during their reproduction season, focusing on trophic resource partitioning.

We analysed the trophic spectrum of 498 specimens of *L. vulgaris* and 238 specimens of *T. cristatus* from a macrophyte-rich, shallow pond (0.2–1 m depth) located in the eastern Romanian Carpathian Mountains, near the town of Piatra Neamț (geographical coordinates: 46° 53' 22" N; 26° 19' 21" E; altitude: 340 m ASL). The stomach contents were collected in two different years (1999 and 2000) in March and April, using the stomach flushing method (Joly, 1987). We used a stereo microscope to analyse the stomach content and to identify prey items to the lowest possible taxonomic level (order, family or genus, in some cases). To determine the importance of each taxonomical prey category in the newts' diet we used the frequency of occurrence (%F) (Fagade and Olaniyan, 1972). In order to estimate the niche breadth, we used the Levins-Hurlbert index (1968, 1978) and niche overlap was quantified through Pianka (1973) and MacArthur-Levins (1967) indices using EcoSim7 Version 7 (Gotelli and Entsminger, 2004).

Food was present in the tracts of 91% of *T. cristatus* and 85% of *L. vulgaris* individuals. The diet of both species was mainly based on isopod crustaceans (Asellidae), freshwater snails (Lymnaeidae) and earthworms (Lumbricidae). The frequency of Lumbricidae in the diet of *T. cristatus* varied greatly between the two study years (from 4.53% in 1999 to 38.46% in 2000) (Table I). The gastrointestinal content of the two species was dominated by representatives of Asellidae family (50.48% in 1999 and 29.92% in 2000 for *T. cristatus*; 50.79% in 1999 and 40.32% in 2000 for *L. vulgaris*) (Table I). *L. vulgaris* fed on a wider variety of food items than *T. cristatus*, although the frequency of occurrence of most of the items was very low (Table I). Previous studies documented the presence of the following invertebrate groups in the gastrointestinal content of *L. vulgaris*: cladocerans, copepods, ostracodes and nematoceran (e.g. David et al., 2009; Covaciu-Marcov et al., 2010). *T. vulgaris* feeds mostly on isopods, nematodes nematoceran and ephemeropteran larvae (e.g. Cicort-Lucaciu et al., 2005; David et al., 2009).

Both *T. cristatus* and *L. vulgaris* had trophic niche breadth values higher than 0.5 (Table II) and the degree of niche overlap (Pianka index) between the two species remained very high (>0.6) during the two years (Table IV). During the spring of 1999 the two species expressed the same high niche overlap upon each other (>0.9), but in 2000 the situation changed, and *L. vulgaris* exercised a higher pressure upon *T. cristatus* (0.75; Table III).

Comparative feeding ecology studies of the two species suggest that *T. cristatus* consumes larger prey than *L. vulgaris* due to a larger body size (Avery, 1968). Nevertheless, our data indicate that even if large body size enables a wider range of prey sizes to be consumed, both large (*T. cristatus*) and small newts (*L. vulgaris*) consume small prey of similar size. Apart from the size differences, our data confirm that both species of newts appear to be opportunistic feeders (Fasola and Canova, 1992; Kutrup et al., 2005), consuming prey which is dominant in the pond and/or is easier to catch (Kopecky et al., 2011; Stephens and Krebs, 1986). Other items which were often found in newts' stomachs are their own skin sloughs which are consumed mainly during the breeding season and this could be a consequence of the physiological shift connected with environmental shift (Kopecky et al., 2011). Aquatic plant fragments

Table I

Frequency of occurrence of different prey items within the diet of *T. cristatus* and *L. vulgaris* during the spring of 1999 and 2000. Legend: nsa – number of stomachs analysed.

Trophic items		<i>Triturus cristatus</i>		<i>Lissotriton vulgaris</i>	
		Year			
Order	Family	1999	2000	1999	2000
Opisthoptera	Lumbricidae	4.53	38.46	2.11	–
Rhynchobdellida	Glossiphoniidae	–	3.84	–	–
Bassomatophora	Lymnaeidae	13.59	38.46	6.87	4.3
Aranea	Araneidae	0.64	–	1.58	1.61
Decapoda	Astacidae	–	–	–	1.07
Isopoda	Asellidae	50.48	29.92	50.79	40.32
Ephemeroptera	Baetidae	2.58	11.53	2.11	–
	Agrionidae	1.29	3.84	4.23	–
Odonata	Coenagrionidae	–	–	1.05	–
Orthoptera	Acrididae	–	–	1.05	–
	Apionidae	0.64	–	–	1.07
	Hydrophilidae	–	7.69	–	2.15
	Dytiscidae	–	–	1.05	–
Coleoptera	Grynidae	–	–	1.05	–
Trichoptera	*	–	–	–	2.15
Lepidoptera	Zygaenidae	–	–	1.05	–
	Ceratopogonidae	0.97	1.92	2.11	1.61
	Chironomidae	–	–	3.17	1.61
	Culicidae	–	–	1.05	2.68
	Dixidae	–	–	1.05	–
	Muscidae	–	–	2.11	3.22
	Ptychopteridae	–	–	1.05	–
	Syrphidae	–	–	–	2.15
Diptera	Tipulidae	–	–	1.05	2.15
Hymenoptera	Ichneumonidae	–	–	1.05	–
Cypriniformes	Cyprinidae	–	3.84	1.05	3.22
Caudata	Salamandridae	–	–	–	1.07
Anura	Ranidae	–	–	2.11	1.07
nsa		52	309	186	189

Table II

Niche breadth of *T. cristatus* and *L. vulgaris* during the spring of 1999 and 2000.

Species	1999	2000
<i>T. cristatus</i>	0.55	0.68
<i>L. vulgaris</i>	0.71	0.83

Table III

Niche overlap (MacArthur-Levins) for *T. cristatus* and *L. vulgaris* during the spring of 1999 and 2000.

Species	1999		2000	
	<i>T. cristatus</i>	<i>L. vulgaris</i>	<i>T. cristatus</i>	<i>L. vulgaris</i>
<i>T. cristatus</i>		0.97*		0.35
<i>L. vulgaris</i>	0.99*		0.75*	

* high overlap between the two trophic niches (>0.5).

Table IV

Niche overlap (Pianka) for *T. cristatus* and *L. vulgaris* during the spring of 1999 and 2000.

Niche overlap	1999	2000
<i>T. cristatus</i> vs. <i>L. vulgaris</i>	0.81*	0.6*

*high overlap between the two trophic niches (>0.5).

were also consumed by both species. Although the presence of these plant fragments in the stomach content of newts is common, these are most probably ingested accidentally. (e.g. Cicort-Lucaciu *et al.* 2005; Covaciu-Marcov *et al.* 2010). Newt eggs and sometimes larvae are usually found in the newts' diet (Cicort-Lucaciu *et al.*, 2005), nevertheless these were not found in our study. *T. cristatus* has been known to consume adult *L. vulgaris* and the incidence of such predation may be high in certain populations (Hagström, 1979). However, we have not recorded such an occurrence in diet.

Both studied newt species may be considered generalists because their niche breadth is higher than 0.5 (MacArthur-Levins, 1967) and their feeding strategy is "active foraging" (Huey and Pianka, 1981). A high level of niche overlap such as the present case represents a high possibility for competition between two species (MacArthur-Levins, 1967; Pianka, 1976). However, niche overlap alone does not indicate competition unless it can be proved that resources are scarce for one or both species (Schoener, 1974; 2009). The high degree of niche overlap between the two species analysed may lead to a temporary competition for the resources or to food partitioning throughout the year according to their necessities.

ACKNOWLEDGEMENTS

This work was supported by a grant of the Romanian National Authority for Scientific Research, CNCS – UEFISCDI, project number PN-II-ID-PCCE-2011-2-0028 (Contract no 4/2012).

REFERENCES

- Alford R.A., Dixon P.M. and Pechmann J.H.K., 2001. Ecology: Global amphibian population declines. *Nature*, 412, 499–500.
- Avery R.A., 1968. Food and feeding relations of three species of *Triturus* (Amphibia Urodela) during the aquatic phases. *Oikos*, 19, 408–412.
- Cicort-Lucaciu A.S., Ardeleanu A., Cupsa D., Naghi N. and Dalea A., 2005. The trophic spectrum of a *Triturus cristatus* (Laurentus 1768) population from Plopiş Mountains area (Bihor County, Romania). *North-West J. Zool.*, 1, 31–39.
- Cogălniceanu D., Aioanei F. and Bogdan M., 2000. Amfibienii din Romania, Determinator. Editura Ars Docenti, Bucureşti.
- Covaciu-Marcov S.D., Cicort-Lucaciu A.S., Mitrea I., Sas I., Caus A.V. and Cupsa D., 2010. Feeding of three syntopic newt species (*Triturus cristatus*, *Mesotriton alpestris* and *Lissotriton vulgaris*) from Western Romania. *North-West J. Zool.*, 6, 95–108.
- David A., Cicort-Lucaciu A.Ş., Roxin M., Pal A. and Nagy-Zachari A.S., 2009. Comparative trophic spectrum of two newts species, *Triturus cristatus* and *Lissotriton vulgaris* from Mehedinţi County, Romania. *Bihorean Biologist.*, 3, 133–137.
- Fagade S.O., Olaniyan C.I., 1972. The biology of the West African shad *Ethmalosa fimbriata* (Bowditch) in the Lagos lagoon, Nigeria. *J. Fish Biol.*, 4, 519–533.
- Fasola M. and Canova L., 1992. Feeding-habits of *Triturus vulgaris*, *Telenomus cristatus* and *T. alpestris* (Amphibia, Urodela) in the Northern Apennines (Italy). *B Zool.*, 59, 273–280.
- Gotelli N.J., Entsminger G.L., 2004. EcoSim: Null models software for ecology. Version 7. Acquired Intelligence Inc&Keseey-Bear. Jericho, VT 05465. <http://geryentsminger.com/ecosim/index.htm>.

- Hagström T., 1979. Population ecology of *Triturus cristatus* and *T. vulgaris* (Urodela) in SW Sweden. *Ecography*, 2, 108–114.
- Hodar J.A., 1996. The use of regression equations for estimation of arthropod biomass in ecological studies. *Acta Oecol.*, 17, 421–433.
- Huey R.B. and Pianka E.R., 1981. Ecological consequences of foraging mode. *Ecology*, 62, 991–999.
- Hurlbert S.H., 1978. The measure of niche overlap and some relatives. *Ecology*, 59, 67–77.
- Iftime A. and Iftime O., 2011. *Triturus cristatus* (Caudata: Salamandridae) feeds upon dead fishes. *Salamandra*, 47, 43–44.
- Joly P., 1987. Le regime alimentaire des Amphibiens méthodes d'étude. *Alytes*, 6, 11–17.
- Kopecky O., Vojar J., Susta F. and Rehak I., 2011. Non-prey items in stomachs of alpine newts (*Mesotriton alpestris*, Laurenti). *Pol. J. Ecol.*, 59, 631–635.
- Kutrup B., Çakir E. and Yilmaz N., 2005. Food of the banded newt, *Triturus vittatus ophryticus* (Berthold, 1846), at different sites in Trabzon. *Turk. J. Zool.*, 29, 83–89.
- Levins R., 1968. Evolution in changing environments: one theoretical explorations. Princeton University Press, NJ.
- MacArthur R. and Levins R., 1967. Limiting similarity convergence and divergence of coexisting species. *Am. Nat.*, 101, 377–385.
- Pianka E.R., 1973. The structure of lizard communities. *Annu. Rev. Ecol. Syst.*, 4, 53–74.
- Pianka E.R., 1976. Competition and niche theory. In: May, R.M. (ed.), Theoretical ecology: principles and applications. Blackwell Scientific Publications, Oxford, 167–196.
- Schabetsberger R. and Jersabek C.D., 1995. Alpine newts (*Triturus alpestris*) as top predators in a high-altitude karst lake - daily food-consumption and impact on the copepod *Arctodiaptomus alpinus*. *Freshwater Biol.*, 33, 47–61.
- Schoener T.W., 1974. Resource partitioning in ecological communities. *Science*, 185, 27–39.
- Schoener T.W., 2009. Ecological niche. In: Levin, S.A. (ed.), The Princeton Guide to Ecology. Princeton University Press, Princeton.
- Semlitsch R.D., 2003. General threats to amphibians. In: Semlitsch, R.D. (ed.), Amphibian Conservation. Smithsonian Institution, Washington DC, 1–7.
- Stephens D.W. and Krebs J.R., 1986. Foraging theory. Princeton University Press, Princeton.
- Stuart S.N., Chanson J.S., Cox N.A., Young B.E., Rodrigues A.S.L., Fischman D.L. and Waller R.W., 2004. Status and trends of amphibian declines and extinctions worldwide. *Science*, 306, 1783–1786.