

# The parasites of the invasive Chinese sleeper *Perccottus glenii* (Fam. Odontobutidae), with the first report of *Nippotaenia mogurndae* in Ukraine

Y. Kvach<sup>(1)\*</sup>, O. Drobiniaik<sup>(1)</sup>, Y. Kutsokon<sup>(2)</sup>, I. Hoch<sup>(3)</sup>

Received February 11, 2013

Revised March 21, 2013

Accepted April 11, 2013

## ABSTRACT

**Key-words:**  
*Chinese sleeper,*  
*parasites,*  
*Nippotaenia*  
*mogurndae,*  
*non-indigenous*  
*species,*  
*neobiota*

The parasites of the Asian invasive fish, Chinese sleeper *Perccottus glenii*, were studied in 6 localities in different parts of Ukraine. In total, 15 taxa of parasites were registered; among them were 1 species of Microsporidia, 5 species of ciliates, 2 species of cestodes, 2 species of trematodes, 2 species of nematodes, 1 species of acanthocephalan, 1 species of parasitic crustacean and 1 mollusk (glochidia). The invasive Chinese sleeper is included as a paratenic host in the life cycle of the parasites of indigenous reptiles in Europe. The non-indigenous cestode *Nippotaenia mogurndae* occurred in the intestine of the Chinese sleeper from the Ivachiv Reservoir (Dniester River basin). This cestode is recorded for Ukrainian fauna for the first time. In addition, 3 species of parasites were recorded in the Chinese sleeper for the first time: *Nicolla skrjabini*, *Cosmocephalus obvelatus* and *Pomphorhynchus laevis*. We note the low similarity among the different localities and the low parasite richness, that suggest that the parasite fauna of the Chinese sleeper in Ukraine is in transition.

## RÉSUMÉ

Les parasites de l'espèce invasive Goujon de l'Amour *Perccottus glenii* (Fam. Odontobutidae) avec le premier signalement de *Nippotaenia mogurndae* en Ukraine

**Mots-clés :**  
*Chinese sleeper,*  
*parasite,*  
*Nippotaenia*  
*mogurndae,*  
*espèces*  
*exotiques,*  
*neobiota*

Les parasites du poisson asiatique invasif, Goujon de l'Amour *Perccottus glenii* (Fam. Odontobutidae), ont été étudiés dans 6 localités de différentes régions d'Ukraine. Au total, 15 taxons de parasites ont été rencontrés, parmi eux, 1 espèce de microsporidies, 5 espèces de ciliés, 2 espèces de cestodes, 2 espèces de trématodes, 2 espèces de nématodes, 1 espèce d'acanthocéphale, 1 espèce de crustacé parasite, et 1 mollusque (glochidies). Le Goujon de l'Amour invasif est considéré comme hôte paraténique dans le cycle de vie des parasites de reptiles indigènes en Europe. Le cestode non-autochtone *Nippotaenia mogurndae* est présent dans l'intestin du Goujon de l'Amour du réservoir d'Ivachiv (bassin de la rivière Dniestr). Ce cestode est rencontré pour la faune ukrainienne pour la première fois. En outre, 3 espèces de parasites ont été enregistrées dans le Goujon de l'Amour pour la première fois : *Nicolla skrjabini*, *Cosmocephalus obvelatus* et *Pomphorhynchus laevis*. Nous notons la faible similitude entre les différentes localités et la richesse parasitaire faible suggérant que la faune parasitaire du Goujon de l'Amour en Ukraine est en transition.

(1) Odessa Branch of the Institute of Biology of the Southern Seas of NAS of Ukraine, Vul. Pushkinska 37, 65011 Odessa, Ukraine

(2) Schmalhausen Institute of Zoology of the NAS of Ukraine, Vul. B. Khmelnytskogo 15, 01601 Kiev, Ukraine

(3) Department of Medical Biology, I.Ya. Gorbachevsky Ternopil State Medical University, Maidan Voli 1, 46001 Ternopil, Ukraine

\* Corresponding author: yuriy.kvach@gmail.com

## INTRODUCTION

The Chinese sleeper *Perccottus glenii* Dybowski, 1877 (Actinopterygii: Odontobutidae) is an invasive fish species indigenous to the freshwaters of Eastern Asia from the Sea of Okhotsk basin in the North, to the Yellow Sea basin in the South (Mori, 1936; Berg, 1949). Now, the invasive Chinese sleeper is widely distributed in the freshwaters of Eastern and Central European countries, such as Belarus, Bulgaria, Estonia, Hungary, Lithuania, Latvia, Moldova, Poland, Romania, Russia, Slovakia, Serbia and Ukraine, where it has high climatic suitability and may continue invasion in the future (Reshetnikov and Ficetola, 2011). In Ukraine, the Chinese sleeper was first found in the upper Dniester River basin in 1980 where it was introduced in the 1970s (Reshetnikov, 2009). It first occurred in the Dnieper river basin near Kiev in 2001, and in the Ros' River (right tributary of the Dnieper River, downstream of Kiev) in 2005 (Sabodash et al., 2002; Kutsokon and Negoda, 2006; Kutsokon, 2010). In the Danube River basin the Chinese sleeper was first recorded in 1995–1996 in the Latorica River, western Ukraine (a part of the western Ukrainian population of the Chinese sleeper), but only in 2011 in the Danube delta (Sivokhop, 1998; Kvach, 2012). This fish is currently found in different parts of the upper streams of the Dniester basin, Transcarpathian waters (Danube basin), in the Dnieper River, and in the Danube River delta.

The parasites of the invasive Chinese sleeper have been thoroughly studied in the European part of Russia (Reshetnikov et al., 2011; Sokolov et al., 2011a, 2011b, 2012) as well as in Poland (Mierzejewska et al., 2010; 2012), including collections from Serbian (Nikolic et al. 2007) and Slovakian waters (Kořuthová et al., 2004, 2008, 2009; Oros and Hanzelová, 2009). In Ukraine, the first study of the Chinese sleeper parasites was carried out by Maslovsky et al. (2011), who reported *Trichodina* sp. and *Crepidostomum farionis* Müller, 1780 in the Vereshchytsia River (Dniester River basin). Davydov et al. (2011) reported parasites of the Chinese sleeper in the Nyvky ponds (Kiev). They reported 8 parasite species: *Saprolegnia* sp., *Trichodina nigra* Lom, 1960, *Trichodina* sp., *Ichthyophthirius multifiliis* Fouquet, 1876, *Bothriocephalus acheilognathi* Yamaguti, 1934, Nematoda sp., *Argulus foliaceus* (Linnaeus, 1758) and Unionidae gen. sp. glochidia.

Three non-indigenous parasites were introduced into European waters together with the Chinese sleeper: the monogenean *Gyrodactylus perccotti* Ergens and Yukhimenko, 1973, the cestode *Nippoetaenia mogurndae* Yamaguti and Miyata, 1940, and the nematode *Philometroides moraveci* Vismanis and Yunchis, 1994 (Kořuthová et al., 2004; Nikolic et al., 2007; Moravec, 2008; Ondračková et al., 2012). The non-indigenous *G. perccotti* is 1 of 2 monogenean species specific to the Chinese sleeper in its native range (Ergens and Yukhimenko, 1973). In Europe it is detected only in the Włocławek Reservoir (Vistula River basin) in Poland; it was not reported from the rivers of the Black Sea basin (Ondračková et al., 2012). *P. moraveci* is a specific parasite of this fish in the Far East (Ermolenko, 2004). It was reported in Serbia as *Philometroides parasiluri* (Yamaguti, 1935) (Nikolic et al., 2007) and later re-identified by Moravec (2008). *N. mogurndae* was first described from the dark sleeper, *Odontobutis obscura* (Temminck and Schlegel, 1845), in Japan (Yamaguti and Miyata, 1940). This tapeworm infects many species of fish of the Odontobutidae family which inhabit the freshwaters of East Asia (Japan, China and Russia) (Yamaguti, 1959). In Europe it was first found in the introduced Chinese sleeper from Slovakia in 2003 (Kořuthová et al., 2004). Later, it was found in the lower Vistula River in 2007 and 2008 (Mierzejewska et al., 2010) and in the middle section of the Volga River basin in 2008–2010 (Reshetnikov et al., 2011).

Information about the parasites of the Chinese sleeper from Ukraine is poor and sporadic, and data about the non-indigenous parasite species are absent. The aim of the present study is to describe the parasite fauna of the Chinese sleeper in Ukrainian waters and to clarify the presence of non-indigenous parasite species in Ukraine.



**Figure 1**

The schematic map of the investigated area. The sampling localities are marked by dots with corresponding numbers. 1. Desna River. 2. Lake Berizka. 3. Novosilky Pond. 4. Trubizh River. 5. Ivachiv Reservoir. 6. Danube delta.

## MATERIAL AND METHODS

Fish were sampled by trawl (8 m long, 0.5 cm in cell diameter) and deep net (1 m/0.5 m, 0.5 cm in cell diameter) at depths from 0.5 to 1.2 m in different water bodies of Ukraine (Figure 1, Table I).

A total of 158 specimens of the Chinese sleeper were studied for parasites. The fish were transported alive in aerated cans to the laboratory of the Odessa Branch of the Institute of Biology of the Southern Seas, stored in aquaria, and dissected within 2 days.

The standard (SL) and total length (TL) of fish were measured before dissection. The skin, fins and gills were scraped off by scalpel, and smears were prepared and studied using light microscopy. If ectoparasites were observed, the smears were dried to prepare the slides. Then, the skin, fins, gills, muscles, brain, eyes, gut, liver, spleen, kidneys, body cavity and mesentery were examined for endoparasites under the dissecting microscope. The encysted larvae were first isolated from cysts and the parasites were fixed in hot formalin. Ciliates were stained with silver nitrate for study of the adhesive disc (Klein, 1958). Trematodes and cestodes were stained in acetic carmine and mounted in Canada balsam for identification (Georgiev *et al.*, 1986). Nematodes and acanthocephalans were studied in glycerine gel temporary mounts. Identification of parasites followed Bauer (1984, 1987), Moravec (1994) and Sudarikov *et al.* (2002), as well as original species descriptions for various taxa.

The parasitological indices were calculated according to Bush *et al.* (1997): prevalence ( $P$  %), intensity (presented as min-max), mean intensity ( $MI$ ) and abundance ( $A$ ). The standard deviation of mean values ( $sd$ ) was calculated. For microparasites, only prevalence was calculated. The gradual scale for the intensity of infection was adopted: 1) sporadic occurrence, “S” – 1 or a few specimens in the examined material; 2) not numerous, “NN” – a few specimens in a few fields of vision; 3) numerous, “N” – up to 20 individuals in most fields of vision; 4) very numerous, “VN” – more than 20 individuals in most fields of vision; 5) mass occurrence, “M” – hundreds of individuals in each field of vision.

**Table I**

Sampling localities, their coordinates, sampling periods, the number of sampled fish in each locality (specimens) and their standard (SL, cm) and total length (TL, cm);  $m \pm sd$  (min-max).

#	Locality name	Locality description	Coordinates	Sampling time	Number of fish sampled in reference period	Total number of sampled fish	Standard length	Total length
1.	Desna River	Near the Lyubychiv Island	50° 45'N 30° 44'E	September 2011	14	29	3.2 ± 0.4 (2.5–4.4)	4.0 ± 0.5 (4.2–5.4)
				August 2012	15			
2.	Lake Berizka	Kiev, on the Hydropark Island, the Dnieper River	50° 26'N 30° 34'E	October 2011	16	31	4.9 ± 1.5 (3.0–11.2)	6.1 ± 1.7 (4.0–13.2)
				April 2012	15			
3.	Novosilky Pond	Kiev, right bank of the Dnieper River	50° 21'N 30° 27'E	November 2011	15	35	5.9 ± 0.9 (3.2–8.7)	7.3 ± 1.7 (4.5–10.8)
				April 2012	20			
4.	Trubizh River	Pereyaslav, Dnieper River basin	50° 4'N 31° 27'E	October 2011	15	30	4.4 ± 1.7 (2.4–10.4)	5.4 ± 2.0 (3.1–12.5)
				May 2012	15			
5.	Ivachiv Reservoir	Near Malashivtsi, the Dniester River basin	49° 40'N 25° 29'E	June 2012	20	20	8.6 ± 5.0 (4.9–27.8)	10.1 ± 5.1 (5.9–29.3)
6.	Danube delta	Vilkove	45° 24'N 29° 35'E	May 2011	2	13	4.6 ± 3.2 (2.5–12.2)	5.6 ± 3.6 (3.2–14.3)
				October 2012	11			

The relative importance of parasites was judged by the core-satellite concept according to their abundance (A):  $>2$  = core species,  $0.6-2$  = secondary species;  $0.2-0.6$  = satellite species; and  $<0.2$  = rare species (Zander *et al.*, 2000).

The Index of Czekanowski-Sørensen, ICS (Sørensen, 1948), was used to compare the parasite faunas:  $ICS = \frac{2c}{a+b} \times 100\%$ , where  $a$  is the number of parasite species found in locality A,  $b$  is the number of parasite species found in locality B, and  $c$  is the number of parasitic species common to both localities. Because of the presence in many cases of unidentified species of *Trichodina*, these ciliates were combined as *Trichodina* spp. for the comparative study.

## RESULTS

A total of 15 taxa of parasites are reported in the Chinese sleeper in Ukraine (Table II); among them were 1 unidentified species of Microsporidia, 5 species of ciliates, 2 species of cestodes, 2 species of trematodes, 2 species of nematodes, 1 species of Acanthocephala, 1 species of parasitic crustacean and 1 species of mollusk glochidia. In some cases the ciliates were not identified to species level but reported as *Trichodina* sp.

Only *Trichodina* spp. were found in all 5 localities studied (Table II). Also, *Holostephanus luhei* Szidat, 1936 metacercariae were found in 2 localities and *Spiroxys contortus* (Rudolphi, 1819) in 3 localities. All other parasites occurred in 1 locality each.

The ciliates were represented by five species: *Trichodina acuta* Lom, 1961, *Trichodina intermedia* Lom, 1960, *Trichodina nigra*, *Trichodina mutabilis* Kazubski and Migala, 1968 and *Trichodina pediculus* Ehrenberg, 1831. *Trichodina* spp. were not identified in sporadic infections. Those sporadic scarce infections were recorded from the Novosilky Pond and the Trubizh

**Table II**

Parasites of the Chinese sleeper in different water bodies of Ukraine.

Parasite species		Desna River	Lake Berizka	Novosilky Pond	Trubizh River	Ivachiv Reservoir	Danube Delta
1		2	3	4	5	6	7
<b>MICROSPORIDIA</b>							
<b>Microsporidia gen. sp.</b>	<i>P</i>	3.4					
	<i>IR</i>	S					
<b>CILIATA</b>							
<b>Trichodina spp.:</b>	<i>P</i>	24.1	51.6	5.7	26.7	75.0	38.5
	<i>IR</i>	S-VN	S-M	S	S-NN	S-M	S-VN
<i>T. acuta</i>			+				
<i>T. intermedia</i>			+				+
<i>T. mutabilis</i>			+				+
<i>T. nigra</i>						+	+
<i>T. pediculus</i>		+	+				+
<i>Trichodina sp.</i>		+		+	+	+	
<b>CESTODA</b>							
<b>Nippotaenia mogurndae</b>	<i>P</i>					55.0	
	<i>MI</i>					5.3 ± 5.3	
	<i>IR</i>					1-19	
	<i>A</i>					2.9	
<b>Ophiotaenia europaea pl</b>	<i>P</i>						7.7
	<i>MI</i>						1.0
	<i>IR</i>						1
	<i>A</i>						0.1
<b>TREMATODA</b>							
<b>Holostephanus luhei met</b>	<i>P</i>	3.4					7.7
	<i>MI</i>	3.0					3.0
	<i>IR</i>	3					3
	<i>A</i>	0.1					0.2
<b>Nicolla skrjabini</b>	<i>P</i>				16.7		
	<i>MI</i>				1.8 ± 0.8		
	<i>IR</i>				1-3		
	<i>A</i>				0.3		
<b>NEMATODA</b>							
<b>Cosmocephalus obvelatus L3</b>	<i>P</i>					10.0	
	<i>MI</i>					2.0 ± 1.4	
	<i>IR</i>					1-3	
	<i>A</i>					0.2	
<b>Spiroxys contortus L3</b>	<i>P</i>	13.8		2.9			7.7
	<i>MI</i>	3.5 ± 2.1		1.0			3.0
	<i>IR</i>	1-6		1			3
	<i>A</i>	0.5		0.03			0.2
<b>ACANTHOCEPHALA</b>							
<b>Pomphorhynchus laevis</b>	<i>P</i>					5.0	
	<i>MI</i>					1.0	
	<i>IR</i>					1	
	<i>A</i>					0.1	

**Table II**  
continued.

Parasite species		Desna River	Lake Berizka	Novosilky Pond	Trubizh River	Ivachiv Reservoir	Danube Delta
		2	3	4	5	6	7
<b>CRUSTACEA</b>							
<i>Argulus foliaceus</i>	<i>P</i>					10.0	
	<i>MI</i>					1.5 ± 0.7	
	<i>IR</i>					1–2	
	<i>A</i>					0.2	
<b>MOLLUSCA</b>							
<i>Pseudoanodonta complanata</i> gl.	<i>P</i>		9.7				
	<i>MI</i>		3.0 ± 1.7				
	<i>IR</i>		2–5				
	<i>A</i>		0.3				

In the table: *P* – prevalence, %; *MI* – mean intensity ± standard deviation; *IR* – intensity range; *A* – abundance; *pl* – plerocercoid; *met* – metacercaria; L3 – 3rd stage larvae; gl. – glochidia.

River (Table II). *T. pediculus* was found in 3 localities: the Desna River, Lake Berizka and the Danube delta. *T. nigra* occurred in 2 localities, Ivachiv Reservoir and the Danube delta. *T. mutabilis* and *T. intermedia* were common in Lake Berizka and the Danube delta. *T. acuta* was only found in Lake Berizka.

Adults of 3 species of helminths (*Nippotaenia mogurndae*, *Nicolla skrjabini* (Iwanitzki, 1928) and *Pomphorhynchus laevis* Müller, 1776) were found. Larvae of 4 other helminth species were also collected: 1 plerocercoid, 1 metacercaria, and 2 3rd stage larvae of nematodes. A plerocercoid of the cestode *Ophiotaenia europaea* and larvae of the nematodes [*Cosmocephalus obvelatus* (Creplin, 1825), *S. contortus*] occurred in the mesentery. The metacercaria of 1 species of trematode (*H. luhei*) was located in the muscles of the stomach.


Only the cestode *N. mogurndae* is considered as a core species (Table II). *N. skrjabini*, *C. obvelatus* L3, *Argulus foliaceus* and *Pseudoanodonta complanata* (Rossmässler, 1835) gl. are considered satellite species. *S. contortus* L3 was a satellite species in the Desna and Danube rivers but rare in the Dnieper River. *H. luhei* was a satellite species in the Desna River but rare in the Danube delta. *O. europaea* was rare.

Among the non-indigenous parasites, only *N. mogurndae* were found in the intestine of the Chinese sleepers from the Ivachiv Reservoir (Dniester basin). Fifty-five percent of fish sampled in the Ivachiv Reservoir were infected with 1–19 parasites (Table II). In addition to scolices with young proglottids, there were numerous detached gravid proglottids. The morphological characteristics of cestodes are typical for this species (Figure 2). The body of cestodes consists of 2–11 proglottids. The anterior 2–4 proglottids are wider than they are long. The scolex is 0.41–0.90 mm long and 0.25–0.50 mm wide. The single apical sucker with a sphincter is round, and 0.18–0.30 mm in diameter. It is located on the top of the scolex.

Gravid proglottids and detached proglottids are elliptical, 1.2–4.3 mm long and 0.8–0.92 mm wide. The testes are anterior and circular, with more than 20 per proglottid. The uterus has transverse coils. The single ovary consists of 2 symmetrical lobes. The vitellaria are placed superiorly, forming 2 reciprocal lobes. Eggs are spherical.

## DISCUSSION

The cestode *Nippotaenia mogurndae* is recorded in Ukraine for the first time. The Chinese sleeper population in the upper Dniester River basin is the oldest in Ukraine, where it was first reported in 1980 (Reshetnikov, 2009). According to Reshetnikov *et al.* (2011), the presence/absence of *N. mogurndae* in the Chinese sleeper population is in agreement with the



**Figure 2**

*Nippotaenia mogurndae* ex.: the Chinese sleeper, *Perccottus glenii*. Locality: Ivachiv Reservoir, Dniester River basin, Ukraine. A – scolex, B – detached gravid proglottid. Scale: 0.2 mm.

vectors of the fish's introduction. The Chinese sleeper population in the Western Ukraine was a source of this fish invasion in the Danube basin in Slovakia, then in other Danube countries (Koščo *et al.*, 2010; Kvach, 2012). The presence of this parasite in Western Ukraine (current data) and in Slovak waters (Kořuthová, 2004; Kořuthová *et al.*, 2008, 2009; Oros and Hanzelová, 2009), as well as its absence in Southern Ukraine, show the vector of the fish invasion. Thus, we predict the occurrence of *N. mogurndae* in the Danube delta in the near future. The presence of this species of cestode in different localities in the Volga basin, but information from the Russian section of the Dnieper basin is absent (Reshetnikov *et al.*, 2011; Sokolov *et al.*, 2012). Because the origins of the Chinese sleeper population in the Ukrainian sector of the Dnieper River are unknown, we cannot consider the invasion vector from Russia to Ukraine.

The first intermediate hosts of *N. mogurndae* are planktonic copepods, such as *Neutrodiaptomus incongruens* (Poppe, 1888), *Eucyclops serrulatus* (Fischer, 1851), *Mesocyclops crassus* (Fischer, 1853) and *M. leuckarti* (Claus, 1857), which are common in Europe (Demshin, 1985; Terek, 1990; Pronin *et al.*, 1998). Kořuthová *et al.* (2008) assume that the increasing occurrence of *N. mogurndae* carries the risk of the potential transfer of this parasite to native fish species with similar ecology. Despite this hypothesis, the fact of *N. mogurndae* occurrence in native fish has not been mentioned until now.

Except for *N. mogurndae*, 2 other helminths used the Chinese sleeper in Ukraine as a definitive host: *Nicola skrjabini* and *Pomphorhynchus laevis*. The latter is detected in the Chinese sleeper for the first time; this is the first record of an acanthocephalan in the invasive population of the Chinese sleeper.

Four helminth species use the Chinese sleeper as a paratenic host. The definitive hosts of 2 of them, *Holostephanus luhei* and *Cosmocephalus obvelatus*, are fish-eating birds. The nematode *C. obvelatus* is recorded in the Chinese sleeper for the first time.

**Table III**

The matrix of the Index of Czekanowski-Sørensen (ICS, %) among the Chinese sleeper parasite fauna in different water bodies of Ukraine (current and published data).

	1	2	3	4	5	6	7	8
1	100	33.33	66.67	18.18	33.33	22.22	33.33	75.00
2		100	50.00	44.44	50.00	28.57	50.00	33.33
3			100	22.22	50.00	28.57	50.00	66.67
4				100	22.22	33.33	22.22	18.18
5					100	28.57	50.00	33.33
6						100	28.57	22.22
7							100	33.33
8								100

1. Desna River (current data), 2. Lake Berizka (current data), 3. Novosilky Pond (current data), 4. Nyvky ponds (Davydov *et al.*, 2011), 5. Trubizh River (current data), 6. Ivachiv Reservoir (current data), 7. Vereshchytsia River (Maslovsky *et al.*, 2010), 8. Danube Delta (current data).

Two other larval parasites use reptiles as definitive hosts. The cestode *Ophiotaenia europaea* is a parasite of the water snakes *Natrix* sp. but the larvae live in Anuran amphibians (Biserkov and Kostadinova, 1997). This tapeworm has already been reported from the Chinese sleeper in the European part of Russia, where it infects from 3.6% to 37.5% of fish (Sokolov *et al.*, 2011b, 2012). The nematode *Spiroxys contortus* is a parasite from the European pond terrapin [*Emys orbicularis* (L., 1758)] but the paratenic hosts are different small-sized fish, insect larvae, tadpoles and adult frogs (Hedrick, 1935; Moravec, 1994). It is considered a common parasite of the invasive Chinese sleeper in Russian waters, infecting up to 100% of fish (Sokolov *et al.*, 2011b, 2012). The invasive Chinese sleeper functions as a paratenic host in the life cycle of the parasites of indigenous reptiles in Europe. The larvae of frogs are important food items of the invasive Chinese sleeper (Reshetnikov, 2001; Plusnina, 2008), therefore they could be a source of larvae of *O. europaea* and *S. contortus*.

The oldest Chinese sleeper population in Ukraine is in the Upper Dniester River basin (Reshetnikov, 2009). This region is characterized by the presence of the specific parasite of the Chinese sleeper, *N. mogurndae*. This is the only core parasite in the parasite community of the Chinese sleeper in Ukraine.

The richness of the Chinese sleeper's parasites in Ukraine is comparable with that of Slovakia, where two species (*Asymphylogdora* sp. and *N. mogurndae*) were found in the Bodrog River, and 3 species [*Orientocreadium gobii* Bychowski & Dubinina, 1954, *N. mogurndae* and *Paraglaridacris gobii* (Szidat, 1938)] in the Latorica River (Kořuthová *et al.*, 2009; Oros and Hanzelová, 2009), in addition to data from some lakes in the Volga River basin: in the lakes Maloe Vetino, Klukvennoe boloto and Ilyevskoe there were 1 to 7 species (Sokolov *et al.* 2012).

Generally, the richness of the Chinese sleeper's parasites in Ukraine is low. For example, 12 species were found in the Neverovski Karyer Reservoir, 13 species in Lake Krugloe (both in the Volga basin), 16 species in Lake Glushitsa Podstepnaya (Don River basin), and 18 species in the Włocławek Reservoir on the Vistula River (Sokolov *et al.*, 2011a, 2011b, 2012; Mierzejewska *et al.*, 2012). The sizes of fish were not larger than in our study: TL = 7.4–21.0 cm in the Neverovski Karyer Reservoir, TL = 4.9–22.8 cm in Lake Krugloe, TL = 5.0–19.5 cm in Lake Glushitsa Podstepnaya, and TL = 4.1–14.4 cm in the Włocławek Reservoir (Sokolov *et al.*, 2011a, 2011b, 2012; Mierzejewska *et al.*, 2012). In all cases the authors did not present data on the average values and standard deviation. Therefore, the data are not statistically comparable; only the disparity of sizes. The Chinese sleeper parasite richness in its native habitats is even higher: 24 species in Lake Sladkoe (Sakhalin Island) and from 11 to 28 species in different localities of the Russian continental Far East (Ermolenko, 2004; Sokolov and Frolov, 2012).

The parasite communities of the Chinese sleeper in different localities of Ukraine are characterized by low similarity (Table III). According to the ICS data, there is relatively high similarity



among the Desna River, the Novosilky Pond and the Danube delta, due to the presence of *S. contortus* and *H. luhei*. Only ciliates, *Trichodina* spp., are common for all localities studied, as in all published reports (Table II; Maslovsky *et al.*, 2010; Davydov *et al.*, 2011). The ciliates are characterized by low host specificity; all *Trichodina* found are generalists, therefore they adapted to new hosts quickly. The minimal similarity is between the Nyvky ponds (Davydov *et al.*, 2011) and Desna River and the Danube River (Table III).

Usually, invasive species have less parasite richness than in their native habitats (Torchin *et al.*, 2003). The release of parasites and pathogens is one of the conditions of successful invasion. This is supported by studies on the invasive gobiid parasites in North America (Kvach and Stepien, 2008). However, long-term observations of invasive gobiid parasites in the same locality (Lake St. Clair) showed increasing parasite richness (Gendron *et al.*, 2012). The parasite fauna of invasive fish in different non-native localities is characterized by low similarity (Kvach and Stepien, 2008; Kvach and Winkler, 2011). The low similarity between different localities shows that the parasite fauna of the Chinese sleeper is in transition. This is suggested by the low parasite richness of the Chinese sleeper in Ukraine, compared with other localities of its new and native habitats.

## ACKNOWLEDGEMENTS

We gratefully acknowledge Prof. Omar M. Amin (Institute of Parasitic Diseases, Scottsdale, AZ, USA) for his help with proofreading the English text.

## REFERENCES

- Bauer O.N. 1984. Opredelitel' parazitov presnovodnyh ryb fauny SSSR [The key to parasites of wresh-water fish of the USSR fauna], Nauka, Leningrad, Vol. 1, 428 p. [in Russian].
- Bauer O.N. 1987. Opredelitel' parazitov presnovodnyh ryb fauny SSSR [The key to parasites of wresh-water fish of the USSR fauna], Nauka, Leningrad, Vol. 3, 583 p. [in Russian].
- Berg L.S., 1949. Ryby presnyh vod SSSR i sopredelnyh stran [Freshwater fishes of the U.S.S.R. and adjacent countries], Vol. 3. Publishing House of Academy of Science of USSR, Moskva-Leningrad, 462 p. [in Russian].
- Biserkov V. and Kostadinova A., 1997. Development of the plerocercoid I of *Ophiotaenia europea* in reptiles. *Int. J. Parasitol.*, 21, 1513–1516.
- Bush A.O., Lafferty K.D., Lotz J.M. and Shostak A.W., 1997. Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *J. Parasitol.*, 83, 575–583.
- Davydov O.N., Kurovskaya L.Ya., Temnikhanov Yu.D. and Neborachek S.I., 2011. Parasites of some invasive fishes of the fresh water. *Gidrobiol. Zhurn.*, 47, 76–89 [in Russian with an English summary].
- Demshin N.I., 1985. Postembryonal development of cestode *Nippotaenia mogurndae* (Nippotaeniidea, Nippotaeniidae). *Parazitologiya*, 19, 39–43 [in Russian with an English summary].
- Ergens R. and Yukhimenko S.S., 1973. Dva novykh vida roda *Gyrodactylus* Nordmann (Monogenoidea) s from *Perccottus glehni* [Two new species of *Gyrodactylus* Nordmann (Monogenoidea) from *Perccottus glehni*]. *Parazitologiya*, 7, 186–188 [in Russian].
- Ermolenko A.V., 2004. Parasite fauna of the Amur sleeper *Perccottus glehni* (Eleotridae). *Parazitologiya*, 38, 257–260 [in Russian with an English summary].
- Gendron A.D., Marcogliese D.J. and Thomas M., 2012. Invasive species are less parasitized than native competitors, but for how long? The case of the round goby in the Great Lakes–St. Lawrence Basin. *Biol. Inv.*, 14, 367–384.
- Georgiev B., Biserkov V. and Genov T., 1986. *In toto* staining method for cestodes with iron acetocarmine. *Helminthol.*, 23, 279–281.
- Hedrick L.R., 1935. The Life History and Morphology of *Spiroxys contortus* (Rudolphi); Nematoda: Spiruridae. *Trans. American Microscop. Soc.*, 54, 307–335.
- Klein B.M., 1958. The dry silver method and its proper use. *J. Protozool.*, 5, 99–103.

- Koščo J., Košuthová L., Košuth P. and Pekárik L., 2010. Non-native fish species in Slovak waters: origins and present status. *Biologia*, 65, 1057–1063.
- Košuthová L., Letková V., Koščo J. and Košuth P., 2004. First record of *Nippotaenia mogurndae* Yamaguti and Miyata, 1940 (Cestoda: Nippotaeniidea), a parasite of *Perccottus glenii* Dybowski, 1877, from Europe. *Helminthol.*, 41, 55–57.
- Košuthová L., Koščo J., Miklisová D., Letková V., Košuth P. and Manko P., 2008. New data on an exotic *Nippotaenia mogurndae* (Cestoda), newly introduced to Europe. *Helminthol.*, 45, 81–85.
- Košuthová L., Koščo J., Letková V., Košuth P. and Manko P., 2009. New records of endoparasitic helminths in alien invasive fishes from the Carpathian region. *Biologia*, 64, 776–780.
- Kutsokon Yu.K., 2010. Distribution and morphological and biological traits of alien fish species in the Ros River basin (tributary to the Dnieper). *Rus. J. Biol. Inv.*, 1, 106–113.
- Kutsokon Yu.K. and Negoda V.V., 2006. Chinese Sleeper, *Perccottus glenii* (Odontobutidae, Perciformes) – a New Species for River Ros' Basin. *Vest. Zool.*, 40, 282 [in Ukrainian with an English title translation].
- Kvach Y., 2012. First record of the Chinese sleeper *Perccottus glenii* Dybowski, 1877 in the Ukrainian part of the Danube delta. *BioInvasions Records*, 1, 25–28.
- Kvach Y. and Stepien C. A., 2008. Metazoan parasites of introduced round and tubenose gobies in the Great Lakes: support for the “enemy release hypothesis”. *J. Great Lakes Res.*, 34, 23–35.
- Kvach Y. and Winkler H.M., 2011. The colonization of the invasive round goby *Neogobius melanostomus* by parasites in new localities in the South-Western Baltic Sea. *Parasitol. Res.*, 109, 769–780.
- Maslovsky O., Tafiychuk R. and Lyesnik V., 2010. Parazytofauna rotana (*Perccottus glenii* Dybowski) u vodoymah zahodu Ukrayiny [Parasite fauna of the Chinese sleeper (*Perccottus glenii* Dybowski) in the water bodies of western Ukraine]. In: Youth and Progress of Biology: abstracts book of the VI International Scientific Conference of Students and Ph.D. Students, Lviv, September 21–24, 2010, 128–129 [in Ukrainian].
- Mierzejewska K., Martyniak A., Kakareko T. and Hliwa P., 2010. First record of *Nippotaenia mogurndae* Yamaguti and Miyata, 1940 (Cestoda, Nippotaeniidae), a parasite introduced with Chinese sleeper to Poland. *Parasitol. Res.*, 106, 451–456.
- Mierzejewska K., Kvach Y., Woźniak M., Kosowska A. and Dziekońska-Rynko J., 2012. Parasites of an Asian fish, the Chinese sleeper *Perccottus glenii*, in the Włocławek Reservoir on the Lower Vistula River, Poland: in search of the key species in the host expansion process. *Comp. Parasitol.*, 79, 23–29.
- Moravec F., 1994. Parasitic nematodes of freshwater fishes of Europe. Academia, Praha, 473 pp.
- Moravec F., 2008. Misidentification of nematodes from the Chinese sleeper *Perccottus glenii* in Europe. *Bull. Europ. Ass. Fish Pathol.*, 28, 86–87.
- Mori T., 1936. Studies on the geographical distribution of freshwater fishes in Eastern Asia. Toppan Print, Tokio, 88 p.
- Nikolic V., Zimonovic P. and Znidarsic T.K., 2007. First record in Europe of a nematode parasite in Amur Sleeper *Perccottus glenii* Dybowski, 1877 (Perciformes: Odontobutidae). *Bull. Eur. Ass. Fish Pathol.*, 27, 36–38.
- Ondračková M., Matějusková I. and Grabowska J., 2012. Introduction of *Gyrodactylus perccotti* (Monogenea) into Europe on its invasive fish host, Amur sleeper (*Perccottus glenii*, Dybowski 1877). *Helminthol.*, 49, 21–26.
- Oros M. and Hanzelová V., 2009. Re-establishment of the fish parasite fauna in the Tisa River system (Slovakia) after a catastrophic pollution event. *Parasitol. Res.*, 104, 1497–1506.
- Plusnina O.V., 2008. Nutrition of Amur sleeper – *Perccottus glenii* Dybowski, 1877 (Odontobutidae, Pisces) in reservoirs of its native and invasion habitats. *Povolzhskiy Ekologicheskii Zhurnal*, 2, 120–125 [in Russian with an English summary].
- Pronin N.M., Selgeby J.H., Litvinov A.G. and Pronina S.V., 1998. Svravnitel'naya ekologiya i parazitofauna ekzoticheskikh vselentsev v Velikiye ozera mira: rotana-goloveshki (*Perccottus glehni*) v oz. Baikal i yersha (*Gymnocephalus cernuus*) v oz. Verkhnee [The comparative ecology and parasite fauna of exotic invaders in the Great Lakes of the world: Amur sleeper (*Perccottus glehni*) in Lake Baikal and ruff (*Gymnocephalus cernuus*) in Lake Superior]. *Sib. J. Ecol.*, 5, 397–406 [in Russian].
- Reshetnikov A.N., 2001. Influence of introduced fish *Perccottus glenii* (Odontobutidae) on amphibians in small waterbodies of Moscow region. *Zhurnal Obshchey Biologii*, 62, 352–361 [in Russian with an English summary].

- Reshetnikov A.N., 2009. The current range of Amur Sleeper *Perccottus glenii* Dybowski, 1877 (Odontobutidae, Pisces) in Eurasia. *Rossiiskii Zhurnal Biologicheskikh Invasii*, 1, 22–35 [in Russian with an English summary].
- Reshetnikov A.N. and Ficetola G.F., 2011. Potential range of the invasive fish rotan (*Perccottus glenii*) in the Holarctic. *Biol. Inv.*, 13, 2967–2980.
- Reshetnikov A.N., Sokolov S.G. and Protasova E.N., 2011. The host-specific parasite *Nippotaenia mogurndae* confirms introduction vectors of the fish *Perccottus glenii* in the Volga river basin. *J. Appl. Ichthyol.*, 27, 1226–1231.
- Sabodash V.M., Tkachenko V.A. and Tsiba A.A., 2002. The *Perccottus glenii* (Pisces, Odontobutidae) population discovered in basins in Kyiv region. *Vest. Zool.*, 36, 90 [in Ukrainian with an English title translation].
- Sivokhop Ya.M., 1998. Pershi znahidky rotanya (*Perccottus glenii*) Dybowski na Zakarpatti [First record of the Chinese sleeper (*Perccottus glenii*) Dybowski in Transcarpathia]. In: Abstracts of the Students Conference (Seria Biology), Patent # 3, Uzhgorod, 44–45 [in Ukrainian].
- Sokolov S.G. and Frolov E.V., 2012. Diversity of parasites in the Amur sleeper (*Perccottus glenii*, Osteichthyes, Odontobutidae) within its native range. *Zool. Zhurn.*, 91, 17–29 [in Russian with an English summary].
- Sokolov S.G., Protasova E.N. and Kholin S.K., 2011a. Parasites of the introduced Amur sleeper, *Perccottus glenii* (Osteichthyes): Alpha-diversity of parasites and age of the host. *Izvestiya RAN, Ser. Biol.*, 5, 584–592 [in Russian with an English summary].
- Sokolov S.G., Protasova E.N., Reshetnikov A.N. and Voropaeva A.L., 2011b. Interactions of the introduced rotan *Perccottus glenii* Dybowski, 1877 (Osteichthyes, Odontobutidae) with aboriginal fish species: a parasitological aspect. *Povolzhskiy Ekologicheskii Zhurnal*, 2, 203–211 [in Russian with an English summary].
- Sokolov S.G., Protasova E.N., Reshetnikov A.N. and Shed'ko M.B., 2012. Parasites of introduced rotan *Perccottus glenii* (Actinopterygii: Odontobutidae) from water bodies of European Russia. *Uspekhi Sovremennoy Biologii*, 132, 477–492 [in Russian with an English summary].
- Sørensen T.A., 1948. A new method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analysis of vegetation on Danish commons. *Kongelige Danske Videnskabernes Selskabs (Biologiske Skrifter)*, 5, 1–34.
- Sudarikov V.E., Shigin A.A., Kurochkin Yu.V., Lomakin V.V., Stenko R.P. and Yurlova N.I., 2002. Metacerkarii trematod – parazity gidrobiontov Rossii [The metacercariae of trematodes – the parasites of gydrobionts in Russia], Nauka, Moscow, Vol. 1, 298 p. [in Russian].
- Terek J., 1990. Net zooplankton of hydromeliorational canals and its relation to selected ecological factors. *Biologia*, 45, 801–814.
- Torchin M.E., Lafferty K.D., Dobson A.P., McKenzie V.J. and Kuris A.M., 2003. Introduced species and their missing parasites. *Nature*, 421, 628–630.
- Yamaguti S., 1959. Systema Helminthum, vol. II. The cestodes of vertebrates. Interscience Publishers, New York, London, 860 p.
- Yamaguti S. and Miyata I., 1940. *Nippotaenia mogurndae* n. sp. (Cestoda) from a Japanese freshwater fish *Mogurnda obscura* (Temm. et Schleg.). *Jap. J. Med. Sci. VI. Bacteriol. Parasitol.*, 1, 213–214.
- Zander C.D., Reimer L.W., Barz K., Dietel G. and Strohbach U., 2000. Parasite communities of the Salzhaff (Northwest Mecklenburg, Baltic Sea). II. Guild communities, with special regard to snails, benthic crustaceans, and small-sized fish. *Parasitol. Res.*, 86, 359–372.